

# BMC2 PCI 2025 Medications Data Dictionary (PCI)

## Blue Cross Blue Shield of Michigan (BMC2)

### PCI Registry

#### Data Collection Definitions

*Definitions updated 12/10/2025*

**Data Abstraction Instructions:** For each medication, record whether it was administered or contraindicated during the specified timeframes.

#### Selections:

- **Contraindicated:** history of allergy, reaction, sensitivity, or intolerance to the medication as documented in the patient's medical record.
- **During:** Medication was administered while the patient was physically in the procedure room.
- **Given:** Medication was administered (use when timeframe is unspecified).
- **Medications at Admission:** medications the patient was taking before the current ASC/non-hospital outpatient visit.
- **Medications at Discharge:** Medications the patient was prescribed or instructed to resume at discharge.
- **Not Given:** Medication was not administered.
- **Post:** Medication was administered after the patient left the procedure room, up to discharge or the next procedure (whichever occurs first).
- **Within 24 hours:** Medication was administered within 24 hours before PCI procedure start and/or during the PCI procedure.

## Combination Therapy

### Data Abstraction Instructions:

For combination drugs, record each individual component unless a specific combination option is available to select.

#### Examples:

- If the patient receives **Zestoretic** (lisinopril + hydrochlorothiazide), record as both **ACE inhibitor** and **diuretic**.
- If the patient receives **Entresto** (sacubitril + valsartan), select "**Entresto**" since it is a specific option.

*Note: Medication lists provided are not all-inclusive.*

## Aldosterone Antagonist

### Data Abstraction Instructions:

At discharge, indicate if an aldosterone antagonist was prescribed or not prescribed.

#### Selections:

- Prescribed
- Not Prescribed

### Supporting Definitions:

Generic	Brand
eplerenone	Inspira
finerenone	Kerendia
spironolactone	Aldactone, Carospir

## Aspirin

Data abstraction instructions: Record if Given, Not Given

#### Selections:

At discharge:

- Given
- Not given
- Contraindicated

- Not Prescribed

#### **Supporting Definitions**

To code "yes" for aspirin, the minimum dose should be at least 75mg.

### **Bempedoic Acid**

#### **Data Abstraction Instructions:**

For Bempedoic acid (including combination drugs), select whether it was prescribed or given.

#### **Selections:**

- Prescribed
- Not Prescribed
- Given
- Not Given

#### **Supporting Definitions:**

Generic	Brand
Bempedoic acid	Nexletol
Bempedoic acid/ezetimibe	Nexlizet

### **Bivalirudin**

#### **Data Abstraction Instructions:**

Record if bivalirudin was administered.

#### **Selections:**

- **Given**  
If "Given," select all timeframes that apply:
  - During (administered in the procedure room)
  - Post (administered after leaving the procedure room, up to discharge or next procedure)
- **Not Given**

#### **Supporting Definitions:**

Generic	Brand
bivalirudin	Angiomax

### **Cangrelor**

#### **Data Abstraction Instructions:**

Record if Cangrelor was administered.

#### **Selections:**

- **Given**  
If "Given," select all timeframes that apply:
  - During
  - Post
- **Not Given**

### **Colchicine**

#### **Data Abstraction Instructions:**

Record whether colchicine (or colchicine/probenecid) was prescribed or given.

#### **Selections:**

- Prescribed
- Not Prescribed
- Given
- Not Given

#### **Supporting Definitions:**

Generic Name	Brand Name
colchicine	Colcrys, Colcigel Gel, Glopberba, Lodoco, Mitigare
colchicine/probenecid	Bememid

## Entresto

### Data Abstraction Instructions:

Record if Entresto (sacubitril and valsartan combination) was prescribed at discharge.

*Do not capture other medications containing angiotensin II receptor blockers. Only record the specific combination of sacubitril and valsartan (Entresto).*

### Selections:

- Prescribed
- Not Prescribed

### Supporting Definitions:

Generic	Brand
sacubitril and valsartan	Entresto

## Eptifibatide (Integritin)

### Data Abstraction Instructions:

Record if eptifibatide was administered during PCI procedures only.

### Selections:

- Given

If "Given," select all timeframes that apply:

- During (administered in the procedure room)
- Post (administered after leaving the procedure room, up to discharge or next procedure)

- Not Given

## GLP-1 Agonist

### Data Abstraction Instructions:

Record if a GLP-1 agonist was prescribed at admission.

### Selections (At Admission):

- Prescribed
- Not Prescribed

### Supporting Definitions:

Generic	Brand Names
albiglutide	Tanzeum
dulaglutide	Trulicity
exenatide	Bydureon BCise, Byetta
liraglutide	Victoza, Saxenda
lixisenatide	Adlyxin
semaglutide	Ozempic, Rybelsus, Wegovy
tirzepatide	Zepbound, Mounjaro

## Hydration

### Data Abstraction Instructions:

Record whether hydration (oral or intravenous) was given or not, and document the volume administered (in mL/cc).

Specify the timing as outlined below.

### Oral (PO) Hydration:

- Post: Record any fluid intake consumed by the patient within **6 hours after** procedure completion.

### Selections:

- Given

- 6hr post: Enter volume in mL or select "N/A" if not documented.

- **Not Given**

**Intravenous (IV) Hydration:**

- **During:** Record the same IV infusion types administered **during** the procedure.
- **Post:** Record these IV fluids administered **within 6 hours after** procedure end.

**Selections:**

- Given
  - During: Enter volume in mL or select "N/A" if not documented.
  - 6hr post: Enter volume in mL or select "N/A" if not documented.
- Not Given

**Range:**

1- 15,000 mL

**IV Vasopressors**

**Data Abstraction Instructions:**

Record if IV vasopressor infusions (not bolus doses) were administered during PCI.

**Selections:**

- Given
 

If "Given," select all agents that apply:

  - Dopamine
  - Norepinephrine
  - Phenylephrine
  - Other (specify)
- Not Given

**Supporting Definitions:**

Agents include:

- Norepinephrine
- Epinephrine
- Dopamine
- Phenylephrine
- Vasopressin
- Dobutamine
- Giapreza (angiotensin II)

**Icosapent Ethyl**

**Data Abstraction Instructions:**

Record if Icosapent ethyl was prescribed at discharge.

**Selections (At Discharge):**

- Prescribed
- Not Prescribed

**Supporting Definitions:**

Generic	Brand
Icosapent Ethyl	Vascepa

**PPI (Proton Pump Inhibitor)**

**Data Abstraction Instructions:**

Record if a proton pump inhibitor (PPI) was prescribed.

**Selections:**

- Prescribed
- Not Prescribed

**Supporting Definitions:**

Generic	Brand Names
dexlansoprazole	Dexilant, Dexilant Solutab
esomeprazole	Nexium, Nexium 24 HR
lansoprazole	Prevacid, Prevacid 24-hour
omeprazole	PriLOSEC, Prilosec OTC
omeprazole/sodium bicarb	Zegerid, Zegerid OTC
pantoprazole	Protonix
rabeprazole	Aciphex, Aciphex Sprinkle
vonoprazan	Voquenza

## Statin

### Data Abstraction Instructions:

Record if a statin was given, not given, or contraindicated.

### Selections:

- Given  
If "Given," select all that apply.
- Not Given
- Contraindicated

### Supporting Definitions:

Generic	Brand Names
atorvastatin	Lipitor, Atorvastatin
fluvastatin	Lescol
lovastatin	Mevacor, Altoprev
pravastatin	Pravachol
simvastatin	Zocor, FloLipid
rosuvastatin	Crestor
pitavastatin	Livalo, Zypatamag
lovastatin + niacin *	Advicor
atorvastatin + amlodipine *	Caduet
simvastatin + ezetimibe *	Vytorin
simvastatin + niacin *	Simcor

\* Combination statin products

## Tirofiban (Aggrastat)

### Data Abstraction Instructions:

Record if tirofiban was administered during PCI procedures only.

### Selections:

- Given  
If "Given," select all that apply:
  - During (administered in the procedure room)
  - Post (administered after leaving the procedure room, up to discharge or next procedure)
- Not Given















